

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

FIG. 10




FIG. 1

✓ 40

42



FIG. 2A

FIG. 2B




FIG. 2B

FIG. 3




FIG. 3

FIG. 4

70

Block Parameters: Gain

Gain

Element-wise gain ( $y = K \cdot u$ ) or matrix gain ( $y = K \cdot u$  or  $y = u \cdot K$ ).

Parameters

Gain:

72

Multiplication: Element-wise ( $K \cdot u$ )

74

☒ Saturate on integer overflow

OK Cancel Help Apply

FIG. 4

20




FIG. 5

2025-09-16 09:16:00




FIG. 6



79




FIG. 7

110

```

Block {
  EvalDialogParams(void) ← 112
  SetupRuntimeParams(void) ← 114
  InternalData: ← 116
    AST[nParams]; ← 118
    NumericDialogParamValues[nParams]; ← 120
    RuntimeParams[nRunTimeParams]; ← 122
  GetAstForDialogParam(int paramIndex)
  SetAstForDialogParam(int paramIndex, ast)
  GetNumericParamValues(int dialogParamIndex)
  SetNumericParamValues(int dialogParamIndex, value)
  GetNumRuntimeParams()
  GetRuntimeParam(int runtimeParamIdx) ← 124
  SetRuntimeParam(int runtimeParamIdx, value)
}

```

FIG. 8

100




FIG. 9




FIG. 10A




FIG. 10B

Figure 1. The effect of the initial concentration of the monomer on the polymerization of  $\alpha$ -methylstyrene initiated by  $\text{TiCl}_4$  in  $\text{CH}_2\text{Cl}_2$  at  $-78^\circ\text{C}$ . The concentration of the initiator was  $0.001\text{ mol/L}$ . The polymerization was terminated by the addition of methanol. The polymerization was carried out in a 100 mL three-necked round-bottomed flask equipped with a magnetic stirrer, thermometer, and nitrogen inlet. The monomer was added to the flask containing the initiator and the solution was stirred for 5 min before the addition of the monomer. The monomer was added to the flask and the solution was stirred for 5 min before the addition of the monomer. The monomer was added to the flask and the solution was stirred for 5 min before the addition of the monomer.





FIG. 11






FIG. 13